



SIGNS OF AN OPIOID OVERDOSE

Know when to take action



Unresponsiveness
or unconsciousness



Slowed or stopped
breathing



Snoring or
rattling sounds



Cold or
clammy skin



Discolored lips
or fingernails

Recovery is possible.

Wisconsin Addiction Recovery Helpline

Call: 211 or 833-944-4673

Text your ZIP code to 898211.

Online: www.addictionhelpwi.org

Visit our website for resources at DoseOfRealityWI.gov



WISCONSIN DEPARTMENT
of HEALTH SERVICES



HOW TO ADMINISTER NASAL NALOXONE



**Reverse an opioid overdose
and prevent death**



Call 911 and follow the operator's instructions. Give rescue breaths, if needed. Continue breaths every 5-6 seconds.



Place tip of naloxone nozzle in either nostril until your fingers touch the nose.



Press the plunger firmly to release the dose of naloxone.



If there is no response after 2-3 minutes, give a second dose. Stay with the person until help arrives.

**You are not alone.
Help is available 24/7.**

Visit our website for resources
at DoseOfRealityWI.gov



WISCONSIN DEPARTMENT
of HEALTH SERVICES





SIGNOS DE UNA SOBREDOSIS DE OPIOIDES

(SIGNS OF AN OPIOID
OVERDOSE)

Cuándo tomar medidas



Ausencia de
respuesta o pérdida
del conocimiento



Respiración lenta
o ausencia de
respiración



Ronquidos o ruidos
vibrantes



Piel fría o pegajosa



Pérdida de color
en labios y uñas

La recuperación es posible.

Línea de ayuda para la recuperación de
adicciones de Wisconsin

Llame al: 211 o al 833-944-4673

Envíe su código postal al 898211.

En línea: www.addictionhelpwi.org

Para acceder a recursos, visite nuestro sitio web en
DoseOfRealityWI.gov.



WISCONSIN DEPARTMENT
of HEALTH SERVICES



CÓMO ADMINISTRAR NALOXONA POR VÍA NASAL



(HOW TO ADMINISTER NASAL NALOXONE)

**Revierta una sobredosis de opioides
y evite la muerte.**



Llame al 911 y siga las instrucciones del operador. Brinde respiraciones de rescate, en caso de que sean necesarias. Continúe brindando las respiraciones cada 5 o 6 segundos.



Coloque la punta de la boquilla de naloxona en cualquiera de las fosas nasales hasta que sus dedos toquen la nariz.



Presione firmemente el émbolo para liberar la dosis de naloxona.



Si la persona no responde después de 2-3 minutos, administre una segunda dosis. Permanezca con la persona hasta que llegue la ayuda.

**No está solo.
Existe ayuda disponible
24/7.**

Para acceder a recursos,
visite nuestro sitio web en
DoseOfRealityWI.gov.



WISCONSIN DEPARTMENT
of HEALTH SERVICES



dose
of
reality